

REPORT DOCUMENTATION PAGE			Form Approved OMB NO. 0704-0188		
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA, 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</p>					
1. REPORT DATE (DD-MM-YYYY) 03-03-2016		2. REPORT TYPE Final Report		3. DATES COVERED (From - To) 5-Aug-2013 - 4-Aug-2014	
4. TITLE AND SUBTITLE Final Report: Investigation of a Neurocognitive Biomarker and of Methods to Mitigate Biases in Cognitive/Perceptual/Emotional Processing			5a. CONTRACT NUMBER W911NF-13-1-0312		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER 611103		
6. AUTHORS Ruth E. Propper, Ph.D.			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAMES AND ADDRESSES Montclair State University Research & Sponsored Programs 1 Normal Avenue, College Hall 309 Montclair, NJ 07043 -1624			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS (ES) U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709-2211			10. SPONSOR/MONITOR'S ACRONYM(S) ARO		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S) 63346-LS-RIP.1		
12. DISTRIBUTION AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision, unless so designated by other documentation.					
14. ABSTRACT Sustained unilateral hand clenching alters perceptual processing and affective/motivational state, presumed to reflect increased hemispheric activity contralateral to the side of motor movement. Data from electroencephalographic and imaging studies are contradictory regarding the relationship between sustained hand clenching and brain activity. To investigate relationships between brain activity, sustained unilateral hand clenching, and changes in affect and perceptual processing, frontal hemispheric activity was measured via Functional Near Infrared Spectroscopy (fNIRS) using derived O2Hb prior to, during, and post sustained unilateral					
15. SUBJECT TERMS Hemispheric activity, Lateralization, Cognition, fNIRS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	15. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON Ruth Propper
a. REPORT UU	b. ABSTRACT UU	c. THIS PAGE UU			19b. TELEPHONE NUMBER 973-655-5201

Report Title

Final Report: Investigation of a Neurocognitive Biomarker and of Methods to Mitigate Biases in Cognitive/Perceptual/Emotional Processing

ABSTRACT

Sustained unilateral hand clenching alters perceptual processing and affective/motivational state, presumed to reflect increased hemispheric activity contralateral to the side of motor movement. Data from electroencephalographic and imaging studies are contradictory regarding the relationship between sustained hand clenching and brain activity. To investigate relationships between brain activity, sustained unilateral hand clenching, and changes in affect and perceptual processing, frontal hemispheric activity was measured via Functional Near-Infrared Spectroscopy (fNIRS), using derived O2Hb prior to, during, and post-sustained unilateral hand clench. Participants' mood and spatial perception were recorded pre- and post-clenching. Sustained unilateral hand clenching altered brain activity and mood, but not spatial perception. O2Hb increased bilaterally following sustained unilateral hand clenching, relative to baseline, regardless of hand. Sustained unilateral hand clenching resulted in greater ipsilateral, compared with contralateral, O2Hb. An interaction between side of hand clench and change in mood was in the direction predicted by theories of hemispheric lateralization of emotion: Following left hand clenching, individuals became more affectively negative, and following right hand clenching, they became more affectively positive. The only relationship between O2Hb and behavioral measures was a positive correlation between left hemisphere O2Hb during hand clenching, and post-clench nervousness.

Enter List of papers submitted or published that acknowledge ARO support from the start of the project to the date of this printing. List the papers, including journal references, in the following categories:

(a) Papers published in peer-reviewed journals (N/A for none)

<u>Received</u>	<u>Paper</u>
-----------------	--------------

TOTAL:

Number of Papers published in peer-reviewed journals:

(b) Papers published in non-peer-reviewed journals (N/A for none)

<u>Received</u>	<u>Paper</u>
-----------------	--------------

TOTAL:

Number of Papers published in non peer-reviewed journals:

(c) Presentations

Number of Presentations: 0.00

Non Peer-Reviewed Conference Proceeding publications (other than abstracts):

Received Paper

TOTAL:

Number of Non Peer-Reviewed Conference Proceeding publications (other than abstracts):

Peer-Reviewed Conference Proceeding publications (other than abstracts):

Received Paper

TOTAL:

Number of Peer-Reviewed Conference Proceeding publications (other than abstracts):

(d) Manuscripts

Received Paper

TOTAL:

Number of Manuscripts:

Books

Received Book

TOTAL:

TOTAL:

Patents Submitted

Patents Awarded

Awards

Graduate Students

<u>NAME</u>	<u>PERCENT SUPPORTED</u>
FTE Equivalent:	
Total Number:	

Names of Post Doctorates

<u>NAME</u>	<u>PERCENT SUPPORTED</u>
FTE Equivalent:	
Total Number:	

Names of Faculty Supported

<u>NAME</u>	<u>PERCENT SUPPORTED</u>
FTE Equivalent:	
Total Number:	

Names of Under Graduate students supported

<u>NAME</u>	<u>PERCENT SUPPORTED</u>
FTE Equivalent:	
Total Number:	

Student Metrics

This section only applies to graduating undergraduates supported by this agreement in this reporting period

The number of undergraduates funded by this agreement who graduated during this period: 0.00

The number of undergraduates funded by this agreement who graduated during this period with a degree in science, mathematics, engineering, or technology fields:..... 0.00

The number of undergraduates funded by your agreement who graduated during this period and will continue to pursue a graduate or Ph.D. degree in science, mathematics, engineering, or technology fields:..... 0.00

Number of graduating undergraduates who achieved a 3.5 GPA to 4.0 (4.0 max scale):..... 0.00

Number of graduating undergraduates funded by a DoD funded Center of Excellence grant for Education, Research and Engineering:..... 0.00

The number of undergraduates funded by your agreement who graduated during this period and intend to work for the Department of Defense 0.00

The number of undergraduates funded by your agreement who graduated during this period and will receive scholarships or fellowships for further studies in science, mathematics, engineering or technology fields: 0.00

Names of Personnel receiving masters degrees

NAME

Total Number:

Names of personnel receiving PHDs

NAME

Total Number:

Names of other research staff

NAME

PERCENT SUPPORTED

FTE Equivalent:

Total Number:

Sub Contractors (DD882)

Inventions (DD882)

Scientific Progress

See Attachment

Technology Transfer

FINAL PROGRESS REPORT (FPR)

Statement of problem studied/ summary of the research projects on which the equipment has been used, including support of the research work described in the proposal: Sustained unilateral hand clenching alters perceptual processing and affective/motivational state, with these alterations presumed to reflect increased hemispheric activity contralateral to the side of motor movement. By increasing the activity of one versus the other hemi-cortex, sustained unilateral hand clenching is thought to result in a processing bias toward the more activated, contralateral, hemisphere, and a concomitant change in behavior (for example, see Beckman, Gröpel, & Ehrlenspiel, 2013; Goldstein, Revivo, Kreitler & Metuki, 2010; Harmon-Jones, 2006; Peterson, Gravens, & Harmon-Jones, 2011; Peterson, Shackman, & Harmon-Jones, 2008; Propper, McGraw, Brunyé, & Weiss, 2013; Schiff & Lamon, 1994; Schiff & Truchon, 1993; Schiff, Guirguis, Kenwood, & Herman, 1998)

Goldstein, Revivo, Kreitler, and Metuki (2010), using a common method for inducing sustained unilateral hand clenching, had participants clench either their right or left hand as hard as possible for 45 seconds, followed by a resting period of 15 seconds, with this sequence of clench-rest being repeated four times. Participants then performed a line bisection task, considered a measure of hemispheric activation and perceptual attention, and the Remote Associates Task (RAT, Mednick, 1968). Sustained unilateral hand clenching of the left hand resulted in increased leftward line bisection bias, suggesting increased right hemisphere activity. Additionally, such clenching resulted in superior RAT performance, with this effect suggested to reflect the increased access to the wider semantic network of the right hemisphere.

Several studies have demonstrated changes in emotional state following sustained unilateral hand clenching. For example, Schiff and Lamon (1994) found that the emotional tone of stories was more negative following left (right hemisphere activation), compared to right (left hemisphere activation), sustained unilateral hand clenching. Similarly, Harmon-Jones (2006) reported increased approach motivational affect following sustained right unilateral hand clenching (left hemisphere activation). The effects of hand clenching on emotional/motivational state are in accord with, and have been interpreted as fitting, theories of cerebral lateralization of emotion/motivation, with the left hemisphere being approach/positive, and the right hemisphere withdrawal/negative (e.g.; Davidson, 2002).

Despite the behavioral evidence suggesting that sustained unilateral hand clenching alters contralateral cerebral function, thereby inducing a processing bias toward the contralateral hemisphere, the neural substrates underlying changes in behavior following sustained motor movements are not clear. Electroencephalographic (EEG) studies of sustained unilateral hand clenching seem to support the hypothesis that these motor movements increase contralateral frontal hemispheric activity, as demonstrated by decreased EEG power in the contralateral hemisphere during clenching (e.g.: Harmon-Jones, 2006; Peterson, Gravens, & Harmon-Jones, 2011; Peterson, Shackman, & Harmon-Jones, 2008). However, examinations of cortical activity using imaging techniques such as Functional Magnetic Resonance Imaging (fMRI) and Functional Near-Infrared Spectroscopy (fNIRS) demonstrate both contralateral and ipsilateral brain activity during or following sustained unilateral hand motor movements (e.g.: Derosière, Alexandre, Bourdillon, Mandrick, Ward, & Perrey, 2013). For example, Shibuya, Kuboyama, and Tanaka (2014), and Shibuya, Sadamoto, Sato, Moriyama, and Iwade (2008) both using fNIRS, demonstrated activation of both contralateral and ipsilateral motor areas during performance of sustained unimanual (finger) force production. Particularly relevant, Derosière et al. (2013),

using fNIRS, examined frontal cortical areas during a sustained unilateral handgrip task. At high levels of force generation, the ipsilateral frontal lobe demonstrated greater activation than that of the contralateral frontal lobe. Given that currently used methods for generating contralateral hemispheric activity to induce changes in behavior require participants to clench their hands ‘as hard as they can’ for a sustained period of time (usually 45 seconds, repeated between 2 to 4 times), and given that stronger clenching is associated with increased ipsilateral relative to contralateral frontal lobe activity (e.g.: Derosi re, Alexandre, Bourdillon, Mandrick, Ward, & Perrey, 2013), the relationship between changes in behavior induced by sustained unilateral hand clenching, and the neuronal processes responsible for such changes following hand clenching, is unclear.

The purpose of the research was to examine motor movement-induced changes in perception and affective/motivational state, and their relationship to brain activity in the prefrontal cortex as measured via fNIRS. Line bisection was chosen because performance has previously been shown to change following sustained unilateral hand clenching (e.g.: Goldstein, Revivo, Kreidler, & Metuki, 2010), with left hand clenching resulting in leftward bias, presumed to indicate increased right hemisphere activity. Emotion was also assessed because changes in mood following sustained unilateral hand clenching have also been reported (e.g.: Harmon-Jones, 2006; Schiff & Lamon, 1994). These emotional changes have been consistent with theories of lateralization of emotional/motivation state (e.g.: Davidson, 2002), such that unilateral clenching of the left hand, presumed to activate the right hemisphere, results in negative or withdrawal motivations, and unilateral clenching of the right hand, presumably via activation of the left hemisphere, results in positive or approach motivations. fNIRS was used to assess prefrontal cortical activity following sustained unilateral hand clenching, and the relationships between brain oxygenation, mood, and line bisection performance were examined. Specifically, In order to investigate the relationship between brain activity, sustained unilateral hand clenching, and changes in affect and perceptual processing, frontal hemispheric activity was measured via Functional Near-Infrared Spectroscopy (fNIRS), using derived O₂Hb prior to, during, and post-sustained unilateral hand clench. Participants’ mood (via the Brief Mood Introspection Scale) and spatial perception (via Line Bisection) were recorded pre- and post-clenching. Additionally, tympanic membrane temperature (TMT), a potential indicator of hemispheric activity (see Propper & Bruny , 2013), was also investigated. Because TMT may indicate hemispheric activity, and therefore has the potential to be a marker for cognitive performance, linking TMT with other neurophysiological markers of hemispheric activity (e.g. fNIRS-derived O₂Hb) would support the utility of this simple measure.

Summary of most important results: Sustained unilateral hand clenching altered brain activity and mood, but not spatial perception. Results revealed increased O₂Hb bilaterally following sustained unilateral hand clenching, relative to baseline, regardless of hand. In agreement with previous fNIRS studies, sustained unilateral hand clenching resulted in greater ipsilateral, compared with contralateral, O₂Hb. (See Appendix A for Figures of relationships between side of unilateral hand clench and O₂Hb.) An interaction between side of hand clench and change in mood was in the direction predicted by theories of hemispheric lateralization of emotion: Following left hand clenching, individuals became more affectively negative, and following right hand clenching, they became more affectively positive. The only relationship between O₂Hb and behavioral measures was a positive correlation between left hemisphere O₂Hb during hand clenching, and post-clench nervousness. No relationship between any measure or manipulation

with TMT was found. Result indicate sustained unilateral hand clenching alters hemispheric activity as well as affective/motivational state. However, the underlying relationship between brain activity and mood is still unclear.

Summary of future research on which the equipment will be used: Future research will investigate variations in muscle strength using EMG in conjunction with behavioral, cognitive and fNIRS monitoring to determine if hand clenching strength is related specifically to changes associated with unilateral hand clenching. Other work will attempt to replicate fNIRS findings of primarily ipsilateral hemispheric activity subsequent to unilateral hand clenching using EEG. Additional research explicitly varying strength of hand clenching (for example, by having participants clench 50% of effort) in examinations using fNIRS and behavioral/cognitive/perceptual measures will further shed light on the interrelationships between hemispheric activity changes as a result of sustained unilateral hand clenching and changes in emotion, cognition, and perception.

BIBLIOGRAPHY

- Beckmann, J., Gröpel, P., & Ehrlenspiel, F. (2013). Preventing motor skill failure through hemisphere-specific priming: Cases from choking under pressure. *Journal of Experimental Psychology: General*, 142(3), 679.
- Davidson, R.J., (2002). Anxiety and affective style: role of prefrontal cortex and amygdala. *Biological Psychiatry*, 51, 68–80.
- Derosiere, G., Alexandre, F., Bourdillon, N., Mandrick, K., Ward, T. E., & Perrey, S. (2014). Similar scaling of contralateral and ipsilateral cortical responses during graded unimanual force generation. *NeuroImage*, 85, 471-477.
- Goldstein, A., Revivo, K., Kreitler, M., & Metuki, N. (2010). Unilateral muscle contractions enhance creative thinking. *Psychonomic Bulletin & Review*, 17(6), 895-899.
- Harmon-Jones, E. (2006). Unilateral right-hand contractions cause contralateral alpha power suppression and approach motivational affective experience. *Psychophysiology*, 43(6), 598-603.
- Peterson, C. K., Gravens, L. C., & Harmon-Jones, E. (2011). Asymmetric frontal cortical activity and negative affective responses to ostracism. *Social Cognitive and Affective Neuroscience*, 6(3), 277-285.
- Peterson, C. K., Shackman, A. J., & Harmon-Jones, E. (2008). The role of asymmetrical frontal cortical activity in aggression. *Psychophysiology*, 45(1), 86-92.
- Propper, R. E., McGraw, S. E., Brunye, T. T., & Weiss, M. (2013). Getting a grip on memory: Unilateral hand clenching alters episodic recall. *PloS One*, 8(4), e62474.
- Schiff, B. B., & Lamon, M. (1994). Inducing emotion by unilateral contraction of hand muscles. *Cortex*, 30(2), 247-254.
- Schiff, B. B., & Truchon, C. (1993). Effect of unilateral contraction of hand muscles on perceiver biases in the perception of chimeric and neutral faces. *Neuropsychologia*, 31(12), 1351-1365.
- Schiff, B. B., Guirguis, M., Kenwood, C., & Herman, C. P. (1998). Asymmetrical hemispheric activation and behavioral persistence: Effects of unilateral muscle contractions. *Neuropsychology*, 12(4), 526.

- Shibuya, K., Kuboyama, N., & Tanaka, J. (2014). Changes in ipsilateral motor cortex activity during a unilateral isometric finger task are dependent on the muscle contraction force. *Physiological Measurement*, 35, 417-428.
- Shibuya, K., Sadamoto, T., Sako, K., Moriyama, M., & Iwadata, M. (2008). Quantification of delayed oxygenation in ipsilateral primary motor cortex compared with contralateral side during a unimanual dominant-hand motor task using near-infrared spectroscopy. *Brain Research*, 1210, 142-147.
- Tanaka, M., & Watanabe, W. (2011). Neural compensation mechanisms to regulate motor output during physical fatigue. *Brain Research*, 1395, 46-52.

APPENDIX A

Fig. 3a-3d fNIRSoft-derived frontal-view topograph, with a 190 threshold for visualization.
'Warmer' colors (i.e.: red, orange, yellow) denote increasing O₂Hb

3a O₂Hb in the left and right hemispheres during 30 seconds of the second left hand clenching condition

Fig. 3b O₂Hb in the left and right hemispheres during the 20 seconds post-clenching in the left hand clenching condition

Fig. 3c O₂Hb in the left and right hemispheres during 30 seconds of the second right hand clenching condition

Fig. 3d O₂Hb in the left and right hemispheres during the 20 seconds post-clenching in the right hand clenching condition

Figure 3a




Figure 3b




Figure 3c



Figure 3d

